

**SupplementaryTable1. The frequency of allergens introduced in skin prick test in nasal polyposis.**

<b>Allergen</b>	<b>Frequencyno (%)</b>	<b>CI95%</b>
Chenopodiaceous	25 (22.1)	14.39-29.86
Russian Thistle	26 (23)	15.16-30.85
Dust Mite	23 (20.4)	12.85-27.86
Rye Grass	31 (27.4)	19.12-35.75
Cock Roach	9 (8)	2.92-13.01
Aspergillus	8 (7.1)	2.30-11.66
Timothy Grass	18 (15.9)	9.11-22.75
Total	113 (100)	

**Supplementary Table 2. The frequency of salicylate contained food allergy in nasal polyposis.**

<b>Salicylate Food</b>	<b>Frequency no(%)</b>	<b>CI 95%</b>
Green Pepper	36 (49.3)	37.65-60.98
Red Grape	16 (21.9)	12.27-31.57
Tomato	15 (20.5)	11.12-29.98
Cinnamon	9 (12.3)	4.66-20.00
Peanut	8 (11)	3.67-18.25
Sago	5 (6.8)	0.96-12.74
Thyme	4 (5.5)	0.17-10.79
Mustard	3 (4.1)	0.00-8.74
Apricot	3 (4.1)	0-8.74
Curry powder	3 (4.1)	0.00-8.74
Drill	2 (2.7)	0-6.56
Cucumber	2 (2.7)	0-6.57
Olive	2 (2.7)	0-6.55
Spinage	2 (2.7)	0-6.55
Razz	2 (2.7)	0-6.57
Cherry	1 (1.4)	0-4.08
Apple	1 (1.4)	0-4.08
Mushroom	0 (0)	00.0-00.0
Broccoli	0 (0)	00.0-00.0
Almond	0 (0)	00.0-00.0
<b>Total</b>	<b>73 (100)</b>	